

PRESCRIBING INFORMATION

ARNIL[®] Tablets
(Diclofenac Tablets B.P)

آرنیل ٹیبلٹس

Composition:

Arnil 50 Tablets:

Each enteric coated tablet contains:

Diclofenac Sodium BP..... 50 mg

Arnil 75 Tablets:

Each enteric coated tablet contains:

Diclofenac Sodium BP..... 75 mg

Description:

Diclofenac Sodium is rapidly absorbed into the plasma with peak levels occurring at nearly two hours. It is extensively bound to proteins. The elimination half life is about 1.2 to 1.8 hours. Diclofenac penetrates rapidly into synovial fluid. Peak levels in the synovial fluid of patients with active rheumatoid arthritis and chronic joints effusions are achieved 2 to 4 hours after peak plasma levels following single or multiple dosing. Higher synovial fluid levels are sustained for upto 12 hours after single or multiple dose therapy.

Indications:

Arnil is indicated for the treatment of pain and inflammation in rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, chronic juvenile arthritis and in other musculo-skeletal disorders including peri-arthritis, bursitis, tendinitis, sprains & strains. Arnil also controls pain and inflammation in acute gout, minor surgery and trauma.

Dosage and Administration:

Arnil 50 Tablets: 1 Tablet T.I.D or as directed by the Physician.

Arnil 75 Tablets: 1 Tablet B.I.D or as directed by the Physician.

Side Effects:

Arnil (Diclofenac Tablets B.P) is generally very well tolerated. Patients may sometimes complain of mild headache, dizziness, nausea, diarrhoea, epigastric pain at the start of treatment which usually disappears within few days without discontinuation of therapy, skin-rash, peripheral oedema and slightly increase in transaminase value has infrequently been reported.

Contraindications:

Sensitivity to Diclofenac Sodium, peptic ulceration, asthma. Where patients are hypersensitive to aspirin or other prostaglandin synthetase inhibitors.

Cautions:

As with other NSAIDs periodic monitoring of liver enzymes, creatinine, blood count and urinalysis should be carried out. Avoid in patients with a history of peptic ulcer, haematemesis or in patients with severe hepatic or renal insufficiency. Rarely as with other NSAIDs, anaphylaxis, laryngeal spasm and bronchospasm has been reported.

Drug Interactions:

Concomitant administration of Diclofenac and Aspirin is not recommended because Diclofenac is displaced from binding sites and biliary excretion is increased, resulting in lower plasma concentrations. Study in healthy subjects indicate that Diclofenac Sodium decreases renal clearance of lithium. Concomitant use of gluco-corticoids with NSAIDs, may aggravate the G.I.side-effects.

Storage:

Store at temperature 15 to 30 °C
away from light & moisture.

Presentation:

Arnil 50 Tablets : Pack of 20 Tablets
Arnil 75 Tablets : Pack of 20 Tablets

خوراک اور طریقہ استعمال :

آرنیل 50 ٹیبلیٹس : ایک گولی دن میں تین مرتبہ

یا ڈاکٹر کی ہدایت کے مطابق استعمال کیجئے

آرنیل 75 ٹیبلیٹس : ایک گولی دن میں دو مرتبہ

یا ڈاکٹر کی ہدایت کے مطابق استعمال کیجئے

اسٹوریج : 15 سے 30 ڈگری سینٹی گریڈ درجہ حرارت پر روشنی اور نمی سے بچا کر رکھئے

brookes

Manufactured by:

Brookes Pharma Private Limited
58 - 59 Sector 15 Korangi Industrial Area
Karachi 74900 Pakistan

