

PRESCRIBING INFORMATION

LEVOCIT[®] 5mg Tablets
(Levocetirizine Dihydrochloride)

لیووسٹ 5 ملی گرام ٹیبلٹس

Composition:

Each film coated tablet contains:
Levocetirizine Dihydrochloride (Brookes Specs)..... 5mg.
Mfg. Specs. Brookes

Pharmacology:

Levocit (Levocetirizine Dihydrochloride) is the R-enantiomer of Cetrizine. It is a piperazine derivative and belongs to non-sedating class of antihistamines. Levocit (Levocetirizine Dihydrochloride) is long acting and has some degree of mast cell stabilizing activity. Levocit is almost free of antimuscarinic activity. Levocit is rapidly absorbed from gastrointestinal tract, it is highly bound to plasma proteins and has long elimination half life of approximately 10 hours. Levocit (Levocetirizine Dihydrochloride) is excreted in urine.

Indications:

- For the symptomatic relief of all Allergic conditions.
- Acute & Chronic persistent Rhinitis.
- Chronic Urticaria.

Dosage & Administration:

The daily recommended dose of Levocit Tablets is 5mg.
In Elderly usual dose is 5mg tablet daily. Dose adjustment is required in patients with compromised renal function.

Children aged 6-12 years: the daily recommended dose is 5mg.

Levocit (Levocetirizine Dihydrochloride) is not recommended below 6 years of age.

Levocit 5mg Tablets are administered orally. Duration of treatment depends on type, duration & course of illness or as advised by the physician.

Dosage in Special Situations:

Patients with impaired Renal function:

Dosage of Levocit is adjusted according to renal function and depends on creatinine clearance rate. Levocit should not be used in end stage renal disease or in patients undergoing dialysis.

Patients with Hepatic impairment:

No dose adjustment is required in impaired hepatic function.

Contra-indications:

Hypersensitivity to the active ingredient or any constituents of the Levocit Tablets.

Side-Effects:

Levocit (Levocetirizine Dihydrochloride) is very well tolerated and has relatively few side effects such as dry mouth, headache and fatigue. Nausea and weight gain are infrequently reported.

Interactions:

Simultaneous administration of Levocit (Levocetirizine Dihydrochloride) with alcohol and CNS depressant drugs may have effects on central nervous system.

Pregnancy & Lactation:

No clinical data is available on use of Levocetirizine Dihydrochloride in Pregnancy & lactation.

Ability to Drive & Use machines:

Extra care should be exercised in vehicle driving and operating machines.

Storage:

Store at temperature 15-30 °C away from light and moisture.

Presentation:

Levocit 5mg Tablets are available in 10 Tablets pack.

خوراک اور طریقہ استعمال:

لیوسٹ کی تجویز کردہ خوراک ایک گولی 5 ملی گرام روزانہ ہے۔

6 سے 12 سال کے بچوں کے لیے:

تجویز کردہ خوراک 5 ملی گرام کی ایک گولی روزانہ ہے۔

6 سال سے کم عمر بچوں کے لیے لیوسٹ تجویز نہیں کی جاتی۔

بزرگ مریضوں میں عمومی خوراک 5 ملی گرام روزانہ ہے۔

اسٹوریج:

15 سے 30 ڈگری سینٹی گریڈ درجہ حرارت پر روشنی اور نمی سے بچا کر رکھیے۔

brookes

Manufactured by:

Brookes Pharma Private Limited
58 - 59 Sector 15 Korangi Industrial Area
Karachi 74900 Pakistan.

206-424

